

Topological complexity of 2-torsion lens spaces and ku -(co)homology

Donald M. Davis

Abstract

We use ku -cohomology to determine lower bounds for the topological complexity of mod- 2^e lens spaces. In the process, we give an almost-complete determination of $ku_*(L^\infty(2^e)) \otimes_{ku_*} ku_*(L^\infty(2^e))$, proving a conjecture of González about the annihilator ideal of the bottom class. Our proof involves an elaborate row reduction of presentation matrices of arbitrary size.

2000 Mathematics Subject Classification: 55M30, 55N15.

Keywords and phrases: Topological complexity, lens space, K -theory.

1 Main Theorems

The determination of the topological complexity of topological spaces has been much studied since its introduction by Farber in [2]. The (normalized) topological complexity, $\overline{\text{TC}}(X)$, of a space X is 1 less than the smallest number of open subsets of $X \times X$ over which the fibration $PX \rightarrow X \times X$, which sends a path σ to $(\sigma(0), \sigma(1))$, has a section. See [4] and [5] for an expanded discussion of this concept, especially as it relates to lens spaces.

Let $L^{2n+1}(t)$ denote the standard $(2n+1)$ -dimensional t -torsion lens space, and let $b(n, e)$, as defined in [5], denote the smallest integer k such that there exists a map

$$(1) \quad L^{2n+1}(2^e) \times L^{2n+1}(2^e) \rightarrow L^{2k+1}(2^e)$$

which when followed into $L^\infty(2^e)$ is homotopic to a restriction of the H -space multiplication of $L^\infty(2^e) = B\mathbb{Z}/2^e$. In [4], it is proved that

$$2b(n, e) \leq \overline{\text{TC}}(L^{2n+1}(2^e)) \leq 2b(n, e) + 1.$$

Thus the following theorem yields a lower bound for $\overline{\text{TC}}(L^{2n+1}(2^e))$. Here and throughout $\alpha(n)$ denotes the number of 1's in the binary expansion of n .

Theorem 1.1. *If $e \geq 2$ and $e \leq \alpha(m) < 2e$, then*

$$b(m + 2^{\alpha(m)-e} - 1, e) \geq 2m - 2^{\alpha(m)-e}.$$

This immediately implies the following result for topological complexity, which might be considered our main result.

Corollary 1.2. *If $e \geq 2$ and $e \leq \alpha(m) < 2e$, then*

$$\overline{\text{TC}}(L^{2m+2^{\alpha(m)-e+1}-1}(2^e)) \geq 4m - 2^{\alpha(m)-e+1}.$$

Other results follow from this and the obvious relation $b(n+1, e) \geq b(n, e)$. The author believes that this result contains all lower bounds for $b(n, e)$ implied by 2-primary connective complex K -theory ku . In [6], a much stronger conjectured lower bound for $b(n, e)$ is given, with the same flavor as our theorem. Their conjecture depends on conjectures about $BP^*(L^{2n+1}(2^e) \times L^{2n+1}(2^e))$, while our theorem depends on a theorem about $ku^*(L^{2n+1}(2^e) \times L^{2n+1}(2^e))$.

Our first new result for topological complexity is

$$\overline{\text{TC}}(L^{2m+7}(2^{\alpha(m)-2})) \geq 4m - 8 \text{ if } \alpha(m) \geq 4.$$

Our theorem is proved by applying $ku^*(-)$ to the map (1), obtaining a contradiction under appropriate choice of parameters. Our main ingredient is the almost-complete determination of $ku^{4n-2d}(L^{2n}(2^e) \times L^{2n}(2^e))$. It is well-known that $ku^* = \mathbb{Z}_{(2)}[u]$ with $|u| = -2$ and that its 2^e -series satisfies

$$[2^e](x) = \sum_{i=1}^{2^e} \binom{2^e}{i} u^{i-1} x^i.$$

It is proved in [3, Proposition 3.1] that

$$(2) \quad ku^{\text{ev}}(L^{2n}(2^e) \times L^{2n}(2^e)) = ku^*[x, y]/(x^{n+1}, y^{n+1}, [2^e](x), [2^e](y)),$$

where $|x| = |y| = 2$. One of our main accomplishments is to give a more useful description of $ku^{4n-2d}(L^{2n}(2^e) \times L^{2n}(2^e))$.

On the other hand, ku -homology, $ku_*(L_{2^e})$, of the infinite-dimensional lens space $L_{2^e} = L^\infty(2^e)$ is the ku_* -module generated by classes z_i , $i \geq 0$, of grading $2i + 1$ with relations

$$\sum_{\ell=0}^i \binom{2^e}{\ell+1} u^\ell z_{i-\ell}, \quad i \geq 0.$$

Here $|u| = 2$ in ku_* . Also, $ku_*(L_{2^e} \times L_{2^e})$ contains $ku_*(L_{2^e}) \otimes_{ku_*} ku_*(L_{2^e})$ as a direct ku_* -summand. We define

$$M_e := ku_*(\Sigma^{-1}L_{2^e}) \otimes_{ku_*} ku_*(\Sigma^{-1}L_{2^e}).$$

It is a ku_* -module on classes $[i, j] := z_i \otimes z_j$ of grading $2i + 2j$, $i, j \geq 0$, with relations

$$(3) \quad \sum_{\ell=0}^i \binom{2^e}{\ell+1} u^\ell [i - \ell, j], \quad i, j \geq 0, \quad \text{and} \quad \sum_{\ell=0}^j \binom{2^e}{\ell+1} u^\ell [i, j - \ell], \quad i, j \geq 0.$$

The desuspending was just for notational convenience. Note that the component of M_e in grading $2d$, which we denote by G_d , is isomorphic to $ku^{4n-2d}(L^{2n}(2^e) \times L^{2n}(2^e))$ under the correspondence

$$u^k [i, j] \leftrightarrow u^k x^{n-i} y^{n-j}.$$

Much of our work goes into an almost-complete description of M_e . The result is described in Section 2.

In [5, Theorem 2.1], it is proved that the ideal

$$I_e := (2^e, 2^{e-1}u, 2^{e-2}u^{3 \cdot 2 - 2}, 2^{e-3}u^{3 \cdot 2^2 - 2}, \dots, 2^1 u^{3 \cdot 2^{e-2} - 2}, u^{3 \cdot 2^{e-1} - 2})$$

annihilates the bottom class $[0, 0]$ of M_e , and in [5, Conjecture 2.1] it is conjectured that I_e is precisely the annihilator ideal of $[0, 0]$ in M_e . One of our main theorems is that this conjecture is true.

Theorem 1.3. *For $e \geq 1$, the annihilator ideal of $[0, 0]$ in M_e is precisely I_e .*

This is immediate from our description of M_e in Section 2. See the remark preceding Theorem 2.2.

After describing M_e in Section 2, we use this description in Section 3 to prove Theorem 1.1. In Section 4, we prove our result for M_6 , and in Section 5, we explain how this proof generalizes to arbitrary M_e .

Finally, in Section 6, we give a different proof of Theorem 1.3 for $e \leq 5$, one which is easily checked by a simple computer verification.

The author wishes to thank Jesús González for suggesting this problem, some guidance as to method, and for providing some computer results which were very helpful for finding a general proof.

2 Description of M_e

Our approach to describing M_e is via an associated matrix P_e of polynomials, which we row reduce. The row-reduced form of P_e is quite complicated, and involves some polynomials which are not completely determined. That is why we call our description “almost complete.” In this section, we approach the description of M_e in three steps.

First we give an introduction to our method, define the polynomial matrices P_e , and give in Table 1, without proof, the reduced form of P_4 , obtained without a computer. The result for P_4 is not used in our general proof, but provides a useful example for comparison. Jesús González obtained an equivalent result using `Mathematica`.

Next we give in Theorem 2.1 an almost-complete description of the reduced form of P_6 . This incorporates all aspects of the general reduced P_e , but is still describable in a moderately tractable way. Finally we give in Theorem 2.2 the general result for P_e , which involves a plethora of indices.

Let G_d denote the component of M_e in grading $2d$. Our ordered set of generators for G_d is

$$(4) \quad [0, d], \dots, [d, 0], u[0, d-1], \dots, u[d-1, 0], \dots, u^d[0, 0].$$

Our final presentation matrix of G_d will be a partitioned matrix

$$\begin{pmatrix} M_{0,0} & M_{0,1} & \dots & M_{0,d} \\ \vdots & \vdots & \vdots & \vdots \\ M_{d,0} & M_{d,1} & \dots & M_{d,d} \end{pmatrix},$$

where $M_{i,j}$ is a $(d+1-i)$ -by- $(d+1-j)$ Toeplitz matrix. The columns in a block $M_{i,j}$ correspond to monomials $u^j[-, -]$.

We will use polynomials to represent the submatrices $M_{i,j}$. A polynomial or power series $p(x) = \alpha_0 + \alpha_1 x + \alpha_2 x^2 + \dots$ corresponds to a Toeplitz matrix (of any size) with $(j+k, j)$ entry equal to α_k . Thus the matrix is

$$\begin{pmatrix} a_0 & 0 & 0 & & \\ \alpha_1 & \alpha_0 & 0 & \ddots & \\ \alpha_2 & \alpha_1 & \alpha_0 & & \\ \alpha_3 & \alpha_2 & \alpha_1 & \ddots & \\ \alpha_4 & \ddots & \ddots & \ddots & \\ \vdots & \vdots & \vdots & \vdots & \end{pmatrix}$$

We define P_e to be the polynomial matrix associated to the partitioned presentation matrix of M_e corresponding to the generators (4) and relations (3). In (11) we depict P_6 .

We let

$$p_n(x) = \frac{x^n - 1}{x - 1} = 1 + x + \cdots + x^{n-1}.$$

We will display a single upper-triangular matrix of polynomials, whose restriction to the first $d+1$ columns yields a presentation of G_d for all d . For example, we will see that the first 8 columns for the reduced form of P_4 are

$$\begin{pmatrix} 16 & 0 & 0 & 4xp_2(x) & 0 & 0 & 0 & 2xp_6(x) \\ & 8 & 0 & 4p_3(x) & 0 & 0 & 0 & 2p_7(x) \\ & & 8 & 0 & 0 & 0 & 0 & 0 \\ & & & 8 & 0 & 0 & 0 & 0 \\ & & & & 4 & 0 & 0 & 0 \\ & & & & & 4 & 0 & 0 \\ & & & & & & 4 & 0 \\ & & & & & & & 4 \end{pmatrix}.$$

This implies that a presentation matrix of G_7 is as below.

$$\begin{pmatrix} 16I_8 & 0 & 0 & M_{0,3} & 0 & 0 & 0 & M_{0,7} \\ & 8I_7 & 0 & M_{1,3} & 0 & 0 & 0 & M_{1,7} \\ & & 8I_6 & 0 & 0 & 0 & 0 & 0 \\ & & & 8I_5 & 0 & 0 & 0 & 0 \\ & & & & 4I_4 & 0 & 0 & 0 \\ & & & & & 4I_3 & 0 & 0 \\ & & & & & & 4I_2 & 0 \\ & & & & & & & 4I_1 \end{pmatrix},$$

where I_t is a t -by- t identity matrix, and

$$M_{0,3} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 4 & 0 & 0 & 0 & 0 \\ 4 & 4 & 0 & 0 & 0 \\ 0 & 4 & 4 & 0 & 0 \\ 0 & 0 & 4 & 4 & 0 \\ 0 & 0 & 0 & 4 & 4 \\ 0 & 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad M_{0,7} = \begin{pmatrix} 0 \\ 2 \\ 2 \\ 2 \\ 2 \\ 2 \\ 2 \\ 0 \end{pmatrix},$$

$$M_{1,3} = \begin{pmatrix} 4 & 0 & 0 & 0 & 0 \\ 4 & 4 & 0 & 0 & 0 \\ 4 & 4 & 4 & 0 & 0 \\ 0 & 4 & 4 & 4 & 0 \\ 0 & 0 & 4 & 4 & 4 \\ 0 & 0 & 0 & 4 & 4 \\ 0 & 0 & 0 & 0 & 4 \end{pmatrix}, \quad M_{1,7} = \begin{pmatrix} 2 \\ 2 \\ 2 \\ 2 \\ 2 \\ 2 \\ 2 \end{pmatrix}.$$

The precise reduced form of P_4 is as in Table 1 below. We do not offer a proof here, but can prove it by the methods of Section 4. We often write p_k instead of $p_k(x)$.

The abelian group that the associated matrix of numbers presents has 276 generators and 276 relations. This associated matrix of numbers is almost, but not quite, in Hermite form. For example, the polynomial in position (2, 17) contains terms such as $2x^5$, and so the associated matrix of numbers will have some 2's sitting far above 2's at the bottom of the column. For the matrix to be Hermite, all nonzero entries above a 2 at the bottom should be 1's. We could obtain such a polynomial in position (2, 17) by subtracting $(x^5 + x^6 + x^9 + x^{10})$ times row 17 from row 2. We have chosen not to do this here because it will be important to our reduction that the first three nonzero entries in column 17 are $\frac{1}{2}p_3(x^4)$ times the corresponding entries of column 9.

By restricting to G_1 , the 8 in position (1, 1) shows that $8u[0, 0] = 0$ in M_4 . Similarly, by restriction to G_4 , the 4 in position (4, 4) implies that $4u^4[0, 0] = 0$. We also obtain $2u^{10}[0, 0] = 0$ and $u^{22}[0, 0] = 0$ from the matrix. The Hermite form of the associated matrix of numbers implies that $8[0, 0]$, $4u^3[0, 0]$, $2u^9[0, 0]$, and $u^{21}[0, 0]$ are all nonzero, and this implies Theorem 1.3 when $e = 4$.

Next we describe the reduced form of P_6 . We let $P_{i,j}$ denote the entry in row i and column j , where the numbering of each starts with 0.

Other than these, the nonzero entries are as described below.

- a. There are none in columns 0–2, 4–6, 10–14, 22–30, 46–62, and 94.
- b. The nonzero entries in columns 3, 7–9, 15–17, 31–33, and 63–65 are as below.

	3	7	8	9	15	16	17
0	$16xp_2$	$8xp_6$			$4xp_{14}$		
1	$16p_3$	$8p_7$			$4p_{15}$		
2			$8x^2p_2(x^2)$	$8xp_6$		$4x^2p_6(x^2)$	$4xp_6p_3(x^4)$
3				$8x^2p_2(x^2)$			$4x^2p_6(x^2)$
4			$8p_3(x^2)$	$8xp_2(x^3)$		$4p_7(x^2)$	$4xp_2(x^3)p_3(x^4)$
5				$8p_3(x^2)$			$4p_7(x^2)$

	31	32	33	63	64	65
0	$2xp_{30}$			xp_{62}		
1	$2p_{31}$			p_{63}		
2		$2x^2p_{14}(x^2)$	$2xp_6p_7(x^4)$		$x^2p_{30}(x^2)$	$xp_6p_{15}(x^4)$
3			$2x^2p_{14}(x^2)$			$x^2p_{30}(x^2)$
4		$2p_{15}(x^2)$	$2xp_2(x^3)p_7(x^4)$		$p_{31}(x^2)$	$xp_2(x^3)p_{15}(x^4)$
5			$2p_{15}(x^2)$			$p_{31}(x^2)$

- c. The nonzero entries in columns 18–21, 34–37, and 66–69 are as in Table 2. Here B refers to everything in the 18–21 block except the $4p_3(x^4)$ -diagonal near the bottom. The \bullet s along a diagonal refer to the entry at the beginning of the diagonal. Each letter q refers to a polynomial. These polynomials are, for the most part, distinct. The meaning of the diagram is that, except for the diagonal near the bottom, each entry in the middle portion equals $\frac{1}{2}p_3(x^8)$ times the corresponding entry in the left portion, and similarly for the right portion, as indicated. More formally, for $18 \leq j \leq 21$ and $i < j - 8$,

$$P_{i,j+16} = \frac{1}{2}p_3(x^8) \cdot P_{i,j} \text{ and } P_{i,j+48} = \frac{1}{4}p_7(x^8) \cdot P_{i,j}.$$

- d. Similarly, the nonzero elements in columns 38 to 45 (other than $P_{i,i}$) are as in Table 3. If C denotes all the entries except the $2p_3(x^8)$ -diagonal near the bottom, then columns 70 to 77 are filled

exactly with $C \cdot \frac{1}{2}p_3(x^{16})$ together with a $p_7(x^8)$ -diagonal going down from (22, 70).

Table 2

	18	19	20	21		34	35	36	37		66	67	68	69
0	0	0	4q	4q										
1	0	0	4q	4q										
2	0	4q	4q	4q										
3	0	0	4q	4q										
4	0	4q	4q	4q										
5	0	0	4q	4q										
6	$4x^4 p_2(x^4)$	0	4q	4q										
7	0	●	0	4q										
8	0	0	●	0										
9	0	0	0	●										
10	$4p_3(x^4)$	0	4q	4q										
11	0	●	0	4q										
12	0	0	●	0										
13	0	0	0	●										

$B \cdot \frac{1}{2}p_3(x^8)$ (rows 34-37)
 $B \cdot \frac{1}{4}p_7(x^8)$ (rows 66-69)
 $2p_7(x^4)$ (diagonal 10-13)
 $p_{15}(x^4)$ (diagonal 11-14)

e. Finally, columns 78 to 93 have a form very similar to Table 3 with q instead of $2q$ and rows going from 0 to 61. The lower two diagonals are $x^{16}p_2(x^{16})$ coming down from (30, 78) and $p_3(x^{16})$ coming down from (46, 78), and these are the only non-leading nonzero entries in column 78.

Table 3

	38	39	40	41	42	43	44	45
0	0	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
1	0	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
2	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
3	0	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
4	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
5	0	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
6	0	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
7	0	0	0	$2q$	$2q$	$2q$	$2q$	$2q$
8	0	0	0	0	$2q$	$2q$	$2q$	$2q$
9	0	0	0	0	0	$2q$	$2q$	$2q$
10	0	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
11	0	0	0	$2q$	$2q$	$2q$	$2q$	$2q$
12	0	0	0	0	$2q$	$2q$	$2q$	$2q$
13	0	0	0	0	0	$2q$	$2q$	$2q$
14	$2x^8 p_2(x^8)$	0	0	0	$2q$	0	$2q$	$2q$
15	0	●	0	0	0	$2q$	0	$2q$
16	0	0	●	0	0	0	$2q$	0
17	0	0	0	●	0	0	0	$2q$
18	0	0	0	0	●	0	0	0
19	0	0	0	0	0	●	0	0
20	0	0	0	0	0	0	●	0
21	0	0	0	0	0	0	0	●
22	$2p_3(x^8)$	0	0	0	$2q$	0	$2q$	$2q$
23	0	●	0	0	0	$2q$	0	$2q$
24	0	0	●	0	0	0	$2q$	0
25	0	0	0	●	0	0	0	$2q$
26	0	0	0	0	●	0	0	0
27	0	0	0	0	0	●	0	0
28	0	0	0	0	0	0	●	0
29	0	0	0	0	0	0	0	●

Now we state the general theorem, of which Theorem 1.3 is an immediate consequence, since the first occurrence of $2^{e-\ell}$ along the diagonal occurs in $(3 \cdot 2^{\ell-1} - 2, 3 \cdot 2^{\ell-1} - 2)$.

Theorem 2.2. *Let $P_{i,j}$ denote the entries in the reduced polynomial matrix for M_e . The nonzero entries are*

i. *For $0 \leq s \leq e-1$ and $3 \cdot 2^s - 2 \leq i < 4 \cdot 2^s - 2$ and $0 \leq t \leq e-1-s$,*

$$P_{i,i+2^{s+1}(2^t-1)} = 2^{e-1-s-t} p_{2^{t+1}-1}(x^{2^s}).$$

ii. *For $0 \leq s \leq e-1$ and $2 \cdot 2^s - 2 \leq i < 3 \cdot 2^s - 2$, $P_{i,i} = 2^{e-s}$ and, for $2 \leq t \leq e-s$,*

$$P_{i,i+2^s(2^t-1)} = 2^{e-s-t} x^{2^s} p_{2^t-2}(x^{2^s}).$$

iii. *For $3 \leq t \leq e$ and $2^t + 2^{t-2} - 2 \leq j \leq 2^t + 2^{t-1} - 3$, there are possibly nonzero entries $P_{i,j} = 2^{e-t} q_{i,j}$ for $0 \leq i < j - 2^{t-1}$, and also, for $1 \leq v \leq e-t$,*

$$P_{i,j+2^t(2^v-1)} = 2^{e-t-v} p_{2^{v+1}-1}(x^{2^{t-1}}) q_{i,j}.$$

This generalizes Table 1 and Theorem 2.1. Note that some of the entries of type ii are among the entries of type iii. Note also that $p_1(x) = 1$, and that in part i for $s = e-1$, we usually just consider the smallest value of i .

3 Proof of Theorem 1.1

In this section, we prove Theorem 1.1 by proving the equivalent statement

(5) if $0 \leq t < e$ and $\alpha(m) = t + e$, then $b(m + 2^t - 1, e) \geq 2m - 2^t$.

The case $t = 0$ is elementary ([5, (1.3)]) and is omitted. We will first prove the following cases of (5) and then will show that all other cases follow by naturality.

Theorem 3.1. *For $1 \leq t < e$,*

(6) $b(3 \cdot 2^{t-1} - 1 + 2^{t+1}B, e) \geq 2^{t+2}B$ if $\alpha(B) = e + t - 1$,

and

(7) $b(2^t - 1 + 2^tB, e) \geq (2B - 1)2^t$ if $\alpha(B) = e + t$.

These are the cases $m = 2^{\alpha(B)-e}(4B + 1)$ and $m = 2^{\alpha(B)-e}B$ of Theorem 1.1 or (5).

Proof. We focus on (6), and then discuss the minor changes required for (7). Let $n = 3 \cdot 2^{t-1} - 1 + 2^{t+1}B$ and suppose there is a map

$$L^{2n}(2^e) \times L^{2n}(2^e) \rightarrow L^{2^{t+3}B-1}(2^e)$$

as in (1). Precompose with the self-map $(1, -1)$ of $L^{2n}(2^e) \times L^{2n}(2^e)$, where -1 is homotopic to the Hopf inverse of the identity. Then, as in [1], we obtain

$$(x - y)^{2^{t+2}B} = 0 \in ku^*(L^{2n}(2^e) \times L^{2n}(2^e)).$$

The result (6) will follow from showing that

$$(x - y)^{2^{t+2}B} \neq 0 \in ku^{2(2n-d)}(L^{2n}(2^e) \times L^{2n}(2^e))$$

with $n = 3 \cdot 2^{t-1} - 1 + 2^{t+1}B$ and $d = 3 \cdot 2^t - 2$. This group is isomorphic to the component group G_d for M_e whose presentation matrix was described in Section 2. The ordered set of generators is obtained as $x^{n-d}y^{n-d}$ multiplied by

$$(8) \quad x^0y^d, \dots, x^dy^0, ux^1y^d, \dots, ux^dy^1, \dots, u^dx^dy^d.$$

We omit the $x^{n-d}y^{n-d}$ throughout our analysis.

One easily shows that

$$\nu \binom{2^{t+2}B}{j} \begin{cases} = \alpha(B) & j = 2^{t+1}B \\ > \alpha(B) & 0 < |2^{t+1}B - j| < 2^{t+1}. \end{cases}$$

Here and throughout $\nu(-)$ denotes the exponent of 2 in an integer. We wish to show that if $t < e$ and $d = 3 \cdot 2^t - 2$, then $2^{e+t-1}x^{d/2}y^{d/2} + 2^{e+t}f(x, y) \neq 0$ in G_d , where $f(x, y)$ is a polynomial of degree d in x and y .

In the reduced matrix for P_e we omit all columns and rows not of the form $3 \cdot 2^i - 3$, $0 \leq i \leq t$. Omitting columns amounts to taking a quotient, and when a column(generator) is omitted the row(relation) with its leading entry can be omitted, too. The resulting matrix is presented below, where the various polynomials q are mostly distinct.

Table 4

	0	3	9	$3 \cdot 2^3 - 3$	\dots	$3 \cdot 2^{t-1} - 3$	$3 \cdot 2^t - 3$
0	2^e	$2^{e-2} p_2$	$2^{e-3} q$	$2^{e-4} q$		$2^{e-t} q$	$2^{e-t-1} q$
3		2^{e-1}	$2^{e-3} p_2(x^2)$	$2^{e-4} q$		$2^{e-t} q$	$2^{e-t-1} q$
9			2^{e-2}	$2^{e-4} p_2(x^4)$		$2^{e-t} q$	$2^{e-t-1} q$
$3 \cdot 2^3 - 3$				2^{e-3}		$2^{e-t} q$	$2^{e-t-1} q$
\vdots					\ddots		
$3 \cdot 2^{t-1} - 3$						2^{e-t+1}	$2^{e-t-1} p_2(x^{2^{t-1}})$
$3 \cdot 2^t - 3$							2^{e-t}

We temporarily ignore the polynomials q and the polynomial $f(x, y)$. The first few relevant relations in the corresponding numerical matrix are $x^{d/2}y^{d/2}$ times the following polynomials. We omit writing powers of u ; they equal the degree of the written polynomial.

$$\begin{aligned}
2^e &+ 2^{e-2}(xy^2 + x^2y) \\
&2^{e-1}xy^2 &+ 2^{e-3}(x^3y^6 + x^5y^4) \\
&2^{e-1}x^2y &+ 2^{e-3}(x^4y^5 + x^6y^3) \\
&&2^{e-2}x^3y^6 &+ 2^{e-4}(x^7y^{14} + x^{11}y^{10}) \\
&&2^{e-2}x^4y^5 &+ 2^{e-4}(x^8y^{13} + x^{12}y^9) \\
&&2^{e-2}x^5y^4 &+ 2^{e-4}(x^9y^{12} + x^{13}y^8) \\
&&2^{e-2}x^6y^3 &+ 2^{e-4}(x^{10}y^{11} + x^{14}y^7).
\end{aligned}$$

From these relations, we obtain

$$\begin{aligned}
(9) \quad 2^{e+t-1} &\sim -2^{e+t-3}(xy^2 + x^2y) \\
&\sim 2^{e+t-5}(x^3y^6 + x^4y^5 + x^5y^4 + x^6y^3) \\
&\sim -2^{e+t-7} \sum_{i=7}^{14} x^i y^{21-i} \\
&\sim \dots \\
&\sim \pm 2^{e-t-1} \sum_{i=2^{t-1}}^{2^{t+1}-2} x^i y^{3 \cdot 2^t - 3 - i} \\
&= \pm 2^{e-t-1} (x^{3 \cdot 2^{t-1}-2} y^{3 \cdot 2^{t-1}-1} + x^{3 \cdot 2^{t-1}-1} y^{3 \cdot 2^{t-1}-2}) \\
&\neq 0,
\end{aligned}$$

since maximum exponents are $3 \cdot 2^{t-1} - 1$. That the last line is nonzero follows from the reduced form of the matrix M_e of relations.

Now we incorporate the polynomials q in the above matrix. We denote by \mathbf{m}_i a monomial or sum of monomials of degree $3 \cdot 2^i - 3$, in x and y . At the first step of the above reduction sequence, we would have

an additional $\sum_{i=2}^t 2^{t+e-i-2} \mathbf{m}_i$. At the second step, we add

$$(10) \quad \sum_{i=3}^t 2^{t+e-i-3} \mathbf{m}'_i.$$

We can incorporate the first monomials for $i \geq 3$ into the second, and we replace $2^{t+e-4} \mathbf{m}_2$ by $\sum_{i \geq 3} 2^{t+e-i-3} \mathbf{m}''_i$ and incorporate these into (10). The third step adds $\sum_{i=4}^t 2^{t+e-i-4} \mathbf{m}'''_i$. We incorporate (10) into this for $i > 3$, while the term in (10) with $i = 3$ is equivalent to a sum which can also be incorporated. Continuing, we end with

$$\sum_{i=t}^t 2^{t+e-i-t} \mathbf{m}_t^{(t)} = 2^{e-t} \mathbf{m}_t^{(t)} = 0,$$

so the q 's contribute nothing.

We easily see that incorporating $2^{e+t} f(x, y)$ also contributes nothing, since

$$2^{e+t} \mathbf{m} \sim 2^{e+t-2} \mathbf{m}_1 \sim 2^{e+t-4} \mathbf{m}_2 \sim \dots \sim 2^{e+t-2t} \mathbf{m}_t = 0.$$

The proof of (7) is very similar. We want to show $(x-y)^{(2B-1)2^t} \neq 0$ in $G_{3,2^t-2}$ if $\alpha(B) = e+t$ and $1 \leq t < e$. For $(B-2)2^t < j < (B+1)2^t$, we have

$$\nu \binom{(2B-1)2^t}{j} \begin{cases} = \alpha(B) - 1 & \text{if } j = (B-1)2^t \text{ or } B \cdot 2^t \\ > \alpha(B) - 1 & \text{other } j. \end{cases}$$

We have factored out $x^{n-d} y^{n-d}$ with $n-d = 2^t B - 2^{t+1} + 1$. Our ordered set of generators is again (8), and our class now, mod higher 2-powers, is $2^{e+t-1}(x^{2^t-1} y^{2^{t+1}-1} + x^{2^{t+1}-1} y^{2^t-1})$. Utilizing the relations similarly to (9), we end with

$$\begin{aligned} & \pm 2^{e+t-1} (x^{2^t-1} y^{2^{t+1}-1} + x^{2^{t+1}-1} y^{2^t-1}) \sum_{i=2^t-1}^{2^{t+1}-2} x^i y^{3 \cdot 2^t - 3 - i} \\ &= \pm 2^{e+t-1} (x^{3 \cdot 2^t - 3} y^{3 \cdot 2^t - 2} + x^{3 \cdot 2^t - 2} y^{3 \cdot 2^t - 3}) \neq 0, \end{aligned}$$

since $x^{d+1} = 0 = y^{d+1}$ (after factoring out $x^{n-d} y^{n-d}$). \square

Proof of (5). The proof is by induction on t . If $t = 1$, the theorem follows from (6) with $m = 4B + 1$ if $m \equiv 1 \pmod{4}$, and from (7) with $m = 2B$ if m is even. If $m \equiv 3 \pmod{4}$, then $\alpha(m + 1) \leq \alpha(m) - 1 = e$, so the result follows from the case $t = 0$ for $n = m + 1$.

Now we assume that the result has been proved for all $t' < t$. If m is odd, then $\alpha(m - 1) = e + t - 1$, so using the induction hypothesis in the middle step,

$$b(m + 2^t - 1, e) \geq b(m - 1 + 2^{t-1} - 1, e) \geq 2(m - 1) - 2^{t-1} \geq 2m - 2^t.$$

If $\nu(m + 2^t) = k$ with $1 \leq k \leq t - 2$, then, noting that $\nu(m) = k$, too,

$$\begin{aligned} \alpha(m - 2^k) &= (t + e) - 2^k + \nu(m \cdots (m - 2^k + 1)) \\ &= t + e - 2^k + (2^k - 1) = t + e - 1. \end{aligned}$$

Therefore

$$b(m + 2^t - 1, e) \geq b(m - 2^k + 2^{t-1} - 1, e) \geq 2(m - 2^k) - 2^{t-1} \geq 2m - 2^t.$$

If $\nu(m) \geq t$, let $m = 2^t B$ with $\alpha(B) = \alpha(m) = t + e$. By (7), we obtain $b(m + 2^t - 1, e) \geq 2m - 2^t$, as desired. If $m = 2^{t-1} + 2^{t+1}B$ with $\alpha(B) = t + e - 1$, then (6) is exactly the desired result.

Finally, if $m = 3 \cdot 2^{t-1} + 2^{t+1}A$ with $\alpha(A) = t + e - 2$, then

$$\begin{aligned} \alpha(m + 2^{t-1}) &= \alpha(A + 1) \\ &= \alpha(A) + 1 - \nu(A + 1) \\ &= e + v \text{ with } v < t. \end{aligned}$$

Thus, by the induction hypothesis,

$$b(m + 2^t - 1, e) \geq b(m + 2^{t-1} + 2^v - 1, e) \geq 2(m + 2^{t-1}) - 2^v \geq 2m - 2^t.$$

□

4 Proof of Theorem 2.1

In this section we prove Theorem 2.1. Because it is a fairly complicated row reduction, we accompany the proof with diagrams of the matrix at several stages of the reduction. Although the proof of Theorem 2.2 in Section 5 is a complete proof and subsumes the much-longer proof for

$e = 6$, we feel that the more explicit example renders the general proof more comprehensible, or perhaps unnecessary.

If M is a Toeplitz matrix corresponding to a polynomial $p(x)$ as described in the preceding section, then the Toeplitz matrix corresponding to the polynomial $(1 + \alpha x + \beta x^2)p(x)$ is obtained from M by adding α times each row to the one below it and β times each row to the row 2 below it. This illustrates how row operations on the matrix of polynomials correspond to row operations on the partitioned matrix of numbers.

Our matrices now refer to the case $e = 6$. The initial partitioned matrix for G_d could be considered as the matrix of numbers associated to the following matrix of polynomials, which has $d + 1$ columns and $2(d + 1)$ rows.

$$(11) \quad \begin{pmatrix} 64 & \binom{64}{2}x & \binom{64}{3}x^2 & \binom{64}{4}x^3 & \cdots \\ 64 & \binom{64}{2} & \binom{64}{3} & \binom{64}{4} & \cdots \\ 0 & 64 & \binom{64}{2}x & \binom{64}{3}x^2 & \cdots \\ 0 & 64 & \binom{64}{2} & \binom{64}{3} & \cdots \\ 0 & 0 & 64 & \binom{64}{2}x & \cdots \\ 0 & 0 & 64 & \binom{64}{2} & \cdots \\ & & \vdots & & \end{pmatrix}$$

The first two row blocks of the associated matrices of numbers have $d + 1$ rows of numbers, the next two d rows, etc., while the sizes of the column blocks are $d + 1, d, \dots$. The first (resp. second) (resp. third) row block corresponds to the first (resp. second) (resp. first) set of relations in (3) with $i + j = d$ (resp. d) (resp. $d - 1$).

Note that if the first two rows and the first column of (11) are deleted, we obtain exactly the initial matrix for G_{d-1} . We may assume that the matrix for G_{d-1} has already been reduced, to Q_{d-1} . Thus we may obtain the reduced form for G_d by taking Q_{d-1} , placing a column of 0's in front of it and the top two rows of (11) above that, and then reducing. By the nature of the matrix (11), the restriction of the reduced form Q_d to its first d columns will be Q_{d-1} .

This is an interesting property. Let Q_d denote the reduced form of the polynomial matrix for G_d . Remove its last column, put a column of 0's in front, put the top two rows of (11) above this, and reduce. The result will be the original matrix, Q_d . We will prove that the matrix described in Theorem 2.1 is correct by removing its last column (the one with the 1 at the bottom), preceding the matrix by a column of 0's and this by the first two rows of (11), and seeing that after reducing, we obtain the original matrix Q_{94} . Because of the initial shifting, each column is determined by the column which precedes it, together with the reduction steps, which justifies the method of starting with the putative answer, shifted. This seems to be a rather remarkable proof. However, the reduction is far from being a simple matter.

Now we describe the steps in the reduction. We begin with the putative answer pushed one unit to the right and two units down, preceded by the first two rows of (11) and a column of 0's. We often write R_i and C_j for row i and column j .

Step 0: Subtract R_0 from R_1 , then divide R_1 by $(1 - x)$, and then subtract xR_1 from R_0 . These rows become

$$\begin{pmatrix} 64 & 0 & -\binom{64}{3}x & -\binom{64}{4}xp_2 & -\binom{64}{5}xp_3 & \cdots & -\binom{64}{64}xp_{62} & 0 & \cdots \\ 0 & \binom{64}{2} & \binom{64}{3}p_2 & \binom{64}{4}p_3 & \binom{64}{5}p_4 & \cdots & \binom{64}{64}p_{63} & 0 & \cdots \end{pmatrix}$$

Divide R_1 by 63, which is the unit part of $\binom{64}{2}$. We now have, in R_0 and R_1 ,

$$P_{i,j} = \begin{cases} 64 & i = j = 0 \\ 32 & i = j = 1 \\ 0 & i + j = 1 \\ u_j 2^{6-\nu(j+1)} xp_{j-1} & i = 0, 2 \leq j \leq 63 \\ u'_j 2^{6-\nu(j+1)} p_j & i = 1, 2 \leq j \leq 63 \\ 0 & 0 \leq i \leq 1, 64 \leq j \leq 94, \end{cases}$$

where u_j is the odd factor of $-\binom{64}{j+1}$, and $u'_j \equiv u_j \pmod{64}$.

Our goal is to reduce this matrix so that the first nonzero entry (which we often call the "leading entry") in R_i is

	13	14	15	16	17	18	19	20	21
0	$32xp_{12}$	$64xp_{13}$	$4xp_{14}$	$64xp_{15}$	$32xp_{16}$	$64xp_{17}$	$16xp_{18}$	$64xp_{19}$	$32xp_{20}$
1	$32p_{13}$	$64p_{14}$	$4p_{15}$	$64p_{16}$	$32p_{17}$	$64p_{18}$	$16p_{19}$	$64p_{20}$	$32p_{21}$
2	0	0	0	$4xp_{14}$	0	0	0	0	$4q$
3	0	0	0	$4p_{15}$	0	0	0	0	$4q$
4	0	0	0	0	$4x^2p_6(x^2)$	$4xp_3(x^4)p_6$	0	$4q$	$4q$
5	0	0	0	0	0	$4x^2p_6(x^2)$	0	0	$4q$
6	0	0	0	0	$4p_7(x^2)$	$4xp_2(x^3)p_3(x^4)$	0	$4q$	$4q$
7	0	0	0	0	0	$4p_7(x^2)$	0	0	$4q$
8	0	0	0	0	0	0	$4x^4p_2(x^4)$	0	$4q$
9	0	0	0	0	0	0	0	$4x^4p_2(x^4)$	0
10	0	0	0	0	0	0	0	0	$4x^4p_2(x^4)$
11	0	0	0	0	0	0	0	0	0
12	0	0	0	0	0	0	$4p_3(x^4)$	0	$4q$
13	0	0	0	0	0	0	0	$4p_3(x^4)$	0
14	8	0	0	0	0	0	0	0	$4p_3(x^4)$
15		8	0	0	0	0	0	0	0
16			8	0	0	0	0	0	0
17				8	0	0	0	0	0
18					8	0	0	0	0
19						8	0	0	0
20							8	0	0
21								8	0
22									8

Although it is just a simple shift, it will be useful to have for reference, in Tables 6 and 7 below, the shifted versions of Tables 2 and 3. These are the relevant portions of the matrix at the outset of the reduction. The shifted version of part b of Theorem 2.1 can be mostly seen in Table 5.

Table 6

	19	20	21	22	35	36	37	38	67	68	69	70
2	0	0	$4q$	$4q$								
3	0	0	$4q$	$4q$								
4	0	$4q$	$4q$	$4q$								
5	0	0	$4q$	$4q$								
6	0	$4q$	$4q$	$4q$								
7	0	0	$4q$	$4q$								
8	$4x^4p_2(x^4)$	0	$4q$	$4q$								
9	0	●	0	$4q$								
10	0	0	●	0								
11	0	0	0	●								
12	$4p_3(x^4)$	0	$4q$	$4q$								
13	0	●	0	$4q$								
14	0	0	●	0								
15	0	0	0	●								

$B \cdot \frac{1}{2}p_3(x^8)$

$B \cdot \frac{1}{4}p_7(x^8)$

$2p_7(x^4)$

$p_{15}(x^4)$

Table 7

	39	40	41	42	43	44	45	46
2	0	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
3	0	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
4	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
5	0	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
6	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
7	0	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
8	0	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
9	0	0	0	$2q$	$2q$	$2q$	$2q$	$2q$
10	0	0	0	0	$2q$	$2q$	$2q$	$2q$
11	0	0	0	0	0	$2q$	$2q$	$2q$
12	0	0	$2q$	$2q$	$2q$	$2q$	$2q$	$2q$
13	0	0	0	$2q$	$2q$	$2q$	$2q$	$2q$
14	0	0	0	0	$2q$	$2q$	$2q$	$2q$
15	0	0	0	0	0	$2q$	$2q$	$2q$
16	$2x^8 p_2(x^8)$	0	0	0	$2q$	0	$2q$	$2q$
17	0	●	0	0	0	$2q$	0	$2q$
18	0	0	●	0	0	0	$2q$	0
19	0	0	0	●	0	0	0	$2q$
20	0	0	0	0	●	0	0	0
21	0	0	0	0	0	●	0	0
22	0	0	0	0	0	0	●	0
23	0	0	0	0	0	0	0	●
24	$2p_3(x^8)$	0	0	0	$2q$	0	$2q$	$2q$
25	0	●	0	0	0	$2q$	0	$2q$
26	0	0	●	0	0	0	$2q$	0
27	0	0	0	●	0	0	0	$2q$
28	0	0	0	0	●	0	0	0
29	0	0	0	0	0	●	0	0
30	0	0	0	0	0	0	●	0
31	0	0	0	0	0	0	0	●

At any stage of the reduction, let \tilde{R}_i denote R_i with its leading entry changed to 0. The first nonzero entry of \tilde{R}_i at the outset occurs

in column

$$(13) \quad \left\{ \begin{array}{ll} i + 5 & 4 \leq i \leq 5 \\ i + 3 & 6 \leq i \leq 7 \\ i + 11 & 8 \leq i \leq 11 \\ i + 7 & 12 \leq i \leq 15 \\ i + 23 & 16 \leq i \leq 23 \\ i + 15 & 24 \leq i \leq 31 \\ i + 47 & 32 \leq i \leq 47 \\ i + 31 & 48 \leq i \leq 63. \end{array} \right.$$

For $i \geq 64$, \tilde{R}_i has no nonzero elements.

The relationship between the three parts of Table 6 and the similar relationship that columns 71 to 78 are mostly $\frac{1}{2}p_3(x^{16})$ times Table 7 will be very important. We call it a “proportionality” relation. We extend it to also include that in rows 4, 5, and 6 we have $C_{18}/C_{10} = \frac{1}{2}p_3(x^4)$, $C_{34}/C_{10} = \frac{1}{4}p_7(x^4)$, and $C_{66}/C_{10} = \frac{1}{8}p_{15}(x^4)$, and similarly in row 4, columns 9, 17, 33, and 65. When we perform row operations involving these rows, these relationships continue to hold. Rows 12–15 and 24–31, where the proportionality relationship does not hold, will not be involved in row operations, since the columns in which their leading entry occurs have all 0’s above the leading entry. (Although rows 0 and 1 are initially nonzero in these columns, clearing out R_0 and R_1 , as in Step 1 below, is a 2-step process, and so \tilde{R}_i for $12 \leq i \leq 15$ or $24 \leq i \leq 31$ will not be combining into R_0 or R_1 , either.)

In Steps 3, 6, 9, and 12, we will divide rows 2, 5, 11, and 23 by x , x^2 , x^4 , and x^8 . It will be important that the entire rows are divisible by these powers of x . We keep track of bounds for the x -divisibility of the unspecified polynomials in Table 6 and 7 and in columns 79 to 94. We postpone this analysis until all the reduction steps have been outlined. Similarly to the proportionality considerations just discussed, divisibility bounds are preserved when we add a multiple of one row to another, in that the x -exponent of $P_{i,j} + cP_{i',j}$ is \geq the minimum of that of $P_{i,j}$ and $P_{i',j}$. The rows, 3, 6–7, 12–15, 24–31, and 48–63, where entries not divisible by x occur will not be used to modify other rows.

Now we begin an attempt to remove most of the binomial coefficients from R_0 and R_1 .

Step 1. The goal is to add multiples of lower rows to R_0 and R_1 to reduce them to

	0	1	2	3	4	5	6	7	...	15	...	31	...	63	...
0	64	0	0	$16xp_2$	0	0	0	$8xp_6$	0	$4xp_{14}$	0	$2xp_{30}$	0	xp_{62}	0
1	0	32	0	$16p_3$	0	0	0	$8p_7$	0	$4p_{15}$	0	$2p_{31}$	0	p_{63}	0

with each 0 referring to all intervening columns. However, we will be forced to bring up some additional entries. We claim that, after Step 1, the nonzero entries $P_{1,j}$, in addition to those in columns $2^t - 1$ listed just above, are combinations of various \tilde{R}_j with $j \geq 5$ and j not in $[6, 8] \cup [12, 16] \cup [24, 32] \cup [48, 64]$. Row 0 is similar but has an extra power of x , since this is true at the outset. Rows 0 and 1 will thus have the requisite proportionality and x -divisibility relations.

It will be useful to note that since at the outset all entries in \tilde{R}_i for $i \geq 2$ are a multiple of $\frac{1}{2}$ times the leading entry at the bottom of their column, then, using (13), $2\tilde{R}_i$ can be killed (reduced to all 0's) by subtracting multiples of lower rows if $i \geq 32$. For example, nonzero entries of \tilde{R}_{32} occur only in C_j with $j \geq 79$. If the entry in $(32, j)$ is a polynomial q , then subtracting qR_{j+1} from $2\tilde{R}_{32}$ kills the entry in C_j without changing anything else, since $\tilde{R}_{j+1} = 0$ for such j .

Similarly $4\tilde{R}_i$ can be killed in two steps if $i \geq 16$, and $8\tilde{R}_i$ can be killed if $i \geq 8$. We can use this observation to kill the entries in R_0 and R_1 in many columns.

For example, if $32 \leq j \leq 46$, then the numerical coefficient in $P_{0,j}$ and $P_{1,j}$ is 0 mod 8, while there is a leading 4 in $(j+1, j)$. Subtracting multiples of $2R_{j+1}$ from R_0 and R_1 kills the entries in $(0, j)$ and $(1, j)$ while bringing up multiples of $2\tilde{R}_{j+1}$. This can be killed by the observation of the previous two paragraphs. This method works to eliminate the entries in R_0 and R_1 in columns 12, 14, 16–18, 20–22, 24–30, and 32–62. (Initial entries in R_0 and R_1 in columns > 63 were all 0.) Since $-\binom{64}{2^t} \equiv 2^{6-t} \pmod{2^{13-2t}}$ for $2 \leq t \leq 6$, the entries in R_0 and R_1 in columns 3, 7, 15, 31, and 63 can be changed to their desired values with pure 2-power coefficients by similar steps.

For C_{23} , we subtract even multiples of R_{24} from R_0 and R_1 to kill the entries. This brings into R_0 and R_1 multiples of 4 in some columns 39 to 46 and even entries in some columns ≥ 71 . The latter entries can be cancelled from below, while cancelling multiples of 4 in C_j for

$39 \leq j \leq 46$ brings up multiples of \tilde{R}_{j+1} . A very similar argument and similar conclusion works for removal of entries in $(0, 19)$ and $(1, 19)$.

Now we consider C_{11} . We subtract multiples of $2R_{12}$ to kill the entries in R_0 and R_1 . This brings up multiples of 8 in C_{19} , C_{21} , and C_{22} , 4 in C_j for $35 \leq j \leq 46$, and 2 in some columns > 64 , the latter of which can be cancelled from below. We kill the earlier elements with multiples of R_{j+1} , leaving a combination of the various \tilde{R}_{j+1}

Column 9 is eliminated similarly, giving multiples of \tilde{R}_{21} , \tilde{R}_{37} , and some others, while columns 8, 10, and 13 are, in a sense, easier since their binomial coefficients are 4 times the number at the bottom of their column, rather than 2. For example, to kill the entry in $(1, 13)$, we first subtract a multiple of $4R_{14}$. This contains a $16q$ in C_{21} , which is killed by a multiple of $2R_{22}$. This brings up a $2q'$ in R_{45} , the killing of which brings up a multiple of \tilde{R}_{46} .

To kill the entry in $(1, 5)$, we subtract a multiple of $2R_6$, which has entries in C_j for many values of $j \geq 9$. We can cancel each of these by subtracting a multiple of R_{j+1} , accounting for the contributions to R_1 of multiples of many \tilde{R}_k with $k \notin [6, 8] \cup [12, 16] \cup [24, 32] \cup [48, 64]$. Killing the entries in C_4 and C_6 is similar.

Finally, to kill the entry in $(1, 2)$, we subtract a multiple of $2R_3$. This brings up entries in columns 4, 8, 16, 32, 64, and others, the killing of which brings up combinations of \tilde{R}_5 , \tilde{R}_9 , \tilde{R}_{17} , and \tilde{R}_{33} , as allowed.

Step 2. Subtract $2R_1$ from R_2 to remove the 64 in $P_{2,1}$. This brings entries into R_2 in columns

$$(14) \quad j \in \{3, 7, 10, 15, 18, 20-22, 31, 34, 36-38, 40-46\}$$

and others with $j \geq 63$. The entry brought into C_j has numerical coefficient equal to $P_{j+1,j}$. These are then killed by subtracting corresponding multiples of R_{j+1} , which brings up into R_2 corresponding multiples of \tilde{R}_{j+1} for j as in (14). From \tilde{R}_4 , this will place $q = 8x^2p_2(x^2)p_3$ in C_9 , $\frac{1}{2}p_3(x^4)q$ in C_{17} , $\frac{1}{4}p_7(x^4)q$ in C_{33} , and $\frac{1}{8}p_{15}(x^4)q$ in C_{65} . This extends the proportionality property of columns 9, 17, 33, 65 to include also row 2.

Now R_2 has $16xp_2$ as its leading entry, in column 4.

Step 3. Divide R_2 by xp_2 . Dividing by a polynomial p of the form $1 + \sum \alpha_i x^i$, such as p_2 , is not a problem. If M is a Toeplitz matrix corresponding to a polynomial q , then the Toeplitz matrix corresponding to q/p is obtained from M by performing the row operations corresponding to finitely many of the terms of the power series $1/p$. Dividing by x is more worrisome, and is the reason for much of our work. In this step it is not a big problem, but later, when we have to divide by x^4 and x^8 , more care is required, which will be handled in Theorem 4.2 after all steps have been described.

We have the important relation

$$(15) \quad p_{2t}/p_2 = p_t(x^2),$$

which implies that the entries in $P_{2,j}$ for $j = 8, 16, 32$, and 64 are now $8p_3(x^2)$, $4p_7(x^2)$, $2p_{15}(x^2)$, and $p_{31}(x^2)$. The relation (15) and its variants will be used frequently without comment. In C_9 , we obtain

$$8x \frac{p_2(x^2)p_3}{p_2} = 8xp_2(x^3) + 16x^3/p_2.$$

We use R_{10} to cancel the second term, at the expense of bringing up multiples of $x^3 \tilde{R}_{10}$ into R_2 . This satisfies proportionality properties, which continue to hold.

Step 4. Subtract $p_3 R_2$ from R_3 to change $P_{3,4}$ to 0. Since

$$(16) \quad p_{2k+1} - p_3 p_k(x^2) = -x^2 p_{k-1}(x^2),$$

we obtain $-8x^2 p_2(x^2)$ in $P_{3,8}$, $-4x^2 p_6(x^2)$ in $P_{3,16}$, and similar expressions in C_{32} and C_{64} . We can change the minus to a plus by adding a multiple of R_9 , R_{17} , etc. This brings up multiples of \tilde{R}_9 , \tilde{R}_{17} , etc., into R_3 , but these maintain proportionality and x -divisibility properties. Note that x -divisibility keeps changing. For example, in Step 3, that of R_2 was decreased by 1, and now all that we can say is that the x -divisibility of R_3 is at least the minimum of that of R_2 and its previous value for R_3 . But this will be handled later.

For the convenience of the reader, we list here columns 0 through 10 at this stage of the reduction. Some of the specific polynomials are not very important, and will later just be called q .

	0	1	2	3	4	5	6	7	8	9	10
0	64	0	0	$16xp_2$	0	0	0	$8xp_6$	0	0	$8x^3p_2p_2(x^3)$
1		32	0	$16p_3$	0	0	0	$8p_7$	0	0	$8x^2p_2(x^2)p_2(x^3)$
2		0	0	0	16	0	0	0	$8p_3(x^2)$	$8xp_2(x^3)$	$8p_3p_3(x^2)$
3			32	0	0	0	0	0	$8x^2p_2(x^2)$	$8xp_6$	$8p_3(x^4)$
4				32	0	0	0	0	0	$8x^2p_2(x^2)$	$8xp_6$
5					32	0	0	0	0	0	$8x^2p_2(x^2)$
6						16	0	0	0	$8p_3(x^2)$	$8xp_2(x^3)$
7							16	0	0	0	$8p_3(x^2)$
8								16	0	0	0
9									16	0	0
10										16	0
11											16

Step 5. Subtract $2R_2$ from R_5 to remove the leading entry in R_5 . If $P_{2,j} = q$ for $j > 4$, then adding qR_{j+1} to R_5 will cancel the subtracted entry, at the expense of adding $q\tilde{R}_{j+1}$ to R_5 . So R_5 gets multiples of \tilde{R}_{j+1} for many values of j in the intervals $[8, 10]$, $[16, 22]$, and $[32, 46]$. The rows that we don't want to bring up are 12–15, 24–31, etc., which contain the lower diagonals in Tables 6 and 7, where neither proportionality nor x -divisibility holds.

Now the leading entry of R_5 is $8x^2p_2(x^2)$ in C_{10} .

Step 6. Divide R_5 by $x^2p_2(x^2)$. We need to know that all entries in R_5 are divisible by x^2 . In Theorem 4.2, we will show that this is true for columns 19–22, 35–46, and 67–94. The only other nonzero entries in R_5 are those in columns 10, 18, 34, and 66 with which it started. See Table 5. The first nonzero entries in R_5 after dividing are 8 in C_{10} and $4p_3(x^4)$ in C_{18} .

Step 7. Subtract multiples of R_5 from rows 0, 1, 2, 3, 4, 6, and 7 to clear out C_{10} in these rows. Because it had been the case that $P_{i,18}/P_{i,10} = \frac{1}{2}p_3(x^4)$ for $0 \leq i \leq 6$, we will now have $P_{i,18} = 0$ for $i \in \{0, 1, 2, 3, 4, 6\}$. Also, by (16), $P_{7,18} = 4(p_7(x^2) - p_3(x^2)p_3(x^4)) = -4x^4p_2(x^4)$. We can change the minus to a plus by adding $x^4p_2(x^4)R_{19}$. Similarly, the only nonzero entries in column 34 (resp. 66) (except for $P_{j+1,j}$) are $2p_7(x^4)$ (resp. $p_{15}(x^4)$) in R_5 , and $2x^4p_6(x^4)$ (resp. $x^4p_{14}(x^4)$) in R_7 . This illustrates why the proportionality relations are important.

Step 8. Subtract $2R_5$ from R_{11} to remove the leading entry in R_{11} . Similarly to Step 5, if $P_{5,j} = q$ for $j > 10$, then adding qR_{j+1} to R_{11} will

cancel the subtracted entry, at the expense of adding $q\tilde{R}_{j+1}$ to R_{11} . So R_{11} gets multiples of \tilde{R}_{j+1} for many values of j in the intervals $[18, 22]$ and $[34, 46]$.

The first 23 columns now are as below.

	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
0	64	0	0	$16xp_2$	0	0	0	$8xp_6$	0	0	0	0	0	0	0
1		32	0	$16p_3$	0	0	0	$8p_7$	0	0	0	0	0	0	0
2			0	0	16	0	0	0	0	$8p_3(x^2)$	0	0	0	0	0
3				32	0	0	0	0	0	$8x^2p_2(x^2)$	0	0	0	0	0
4					32	0	0	0	0	0	$8x^2p_2(x^2)$	0	0	0	0
5						0	0	0	0	0	0	8	0	0	0
6							16	0	0	0	$8p_3(x^2)$	0	0	0	0
7								16	0	0	0	0	0	0	0
8									16	0	0	0	0	0	0
9										16	0	0	0	0	0
10											16	0	0	0	0
11												8	0	0	0
12													8	0	0
13														8	0
14															8
15															8
	15	16	17	18	19	20	21	22							
0	$4xp_{14}$	0	0	0	0	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
1	$4p_{15}$	0	0	0	0	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
2	0	$4p_7(x^2)$	$8q_0p_3(x^4)$	0	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
3	0	$4x^2p_6(x^2)$	$8q_1p_3(x^4)$	0	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
4	0	0	$4x^2p_6(x^2)$	0	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
5	0	0	0	$4p_3(x^4)$	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
6	0	0	$4p_7(x^2)$	0	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
7	0	0	0	$4x^4p_2(x^4)$	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
8	0	0	0	0	$4x^4p_2(x^4)$	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
9	0	0	0	0	0	$4x^4p_2(x^4)$	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
10	0	0	0	0	0	0	$4x^4p_2(x^4)$	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
11	0	0	0	0	0	0	0	$4x^4p_2(x^4)$	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
12	0	0	0	0	0	$4p_3(x^4)$	0	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
13	0	0	0	0	0	0	0	$4p_3(x^4)$	0	$4q$	$4q$	$4q$	$4q$	$4q$	$4q$
14	0	0	0	0	0	0	0	0	$4p_3(x^4)$	0	$4q$	$4q$	$4q$	$4q$	$4q$
15	0	0	0	0	0	0	0	0	0	0	$4p_3(x^4)$	0	$4q$	$4q$	$4q$
16	8	0	0	0	0	0	0	0	0	0	0	0	0	0	0
17		8	0	0	0	0	0	0	0	0	0	0	0	0	0
18			8	0	0	0	0	0	0	0	0	0	0	0	0
19				8	0	0	0	0	0	0	0	0	0	0	0
20					8	0	0	0	0	0	0	0	0	0	0
21						8	0	0	0	0	0	0	0	0	0
22							8	0	0	0	0	0	0	0	0
23								8	0	0	0	0	0	0	0

In addition, we have, at this stage of the reduction:

- 4 in $P_{j+1,j}$ for $23 \leq j \leq 46$, and 2 in $P_{j+1,j}$ for $47 \leq j \leq 94$. Other than that:
- 0 in columns 23 to 30 and 47 to 62.
- A pattern resembling that of columns 15 to 18 in columns 31 to 34 and 63 to 66.
- Columns 35 to 38 (resp. 67 to 70) are $\frac{1}{2}p_3(x^8)$ (resp. $\frac{1}{4}p_7(x^8)$) times columns 19 to 22, except that corresponding to the $4p_3(x^4)$ in rows 12 to 15 we have $2p_7(x^4)$ (resp. $p_{15}(x^4)$).

- e. Columns 39 to 46 resemble Table 7. Columns 71 to 78 are $\frac{1}{2}p_3(x^{16})$ times these, except for the diagonal near the bottom, which is $p_7(x^8)$.
- f. Columns 79 to 94 have a form similar to that of columns 39 to 46.
- g. The x -divisibility in columns 19–22, 39–46, and 79–94 will be described in Theorem 4.2 and its proof.

Step 9. Divide R_{11} by $x^4p_2(x^4)$. We will show in Theorem 4.2 that all entries in R_{11} are divisible by x^4 . The leading entry in row 11 is now a 4 in C_{22} .

Step 10. Subtract multiples of R_{11} from rows 0 to 10 and 12 to 15 to clear out their entries in C_{22} . Similarly to Step 7, we now have that $P_{i,38} = 0$ except for $P_{11,38} = 2p_3(x^8)$, $P_{15,38} = 2x^8p_2(x^8)$, and $P_{39,38} = 4$, with a similar situation in C_{70} . In particular, $P_{15,70} = x^8p_6(x^8) = \frac{1}{2}p_3(x^{16})P_{15,38}$.

Step 11. Subtract $2R_{11}$ from R_{23} , and, similarly to Steps 5 and 8, kill entries subtracted from $P_{23,j}$ for $j > 22$ by adding multiples of R_{j+1} , thus bringing up these multiples of \tilde{R}_{j+1} . The smallest such j is 38, due to the entry in (11, 38) described in the previous step.

Step 12. Now the leading entry of R_{23} is $2x^8p_2(x^8)$ in C_{46} . (This can be seen using (13) and that there have been no other changes to R_{23} in columns less than 62.) Divide R_{23} by $x^8p_2(x^8)$. We will show later that all entries in R_{23} are divisible by x^8 at this stage.

Step 13. Subtract multiples of R_{23} from rows 0 to 22 and 24 to 31 to make their entries in C_{46} equal to 0. Similarly to Step 10, this will cause $P_{i,78} = 0$ except for $P_{23,78} = p_3(x^{16})$, $P_{31,78} = x^{16}p_2(x^{16})$, and $P_{79,78} = 2$.

Step 14. Subtract $2R_{23}$ from R_{47} . This will add multiples of 2 to R_{47} in some columns $j \geq 78$. These can be removed, without any other effect, by subtracting a multiple of R_{j+1} . Now R_{47} has leading entry $x^{16}p_2(x^{16})$ in C_{94} . Divide R_{47} by $x^{16}p_2(x^{16})$, and then subtract multiples of R_{47} from the others to clear out C_{94} .

Step 15. We are now in the situation described in the paragraph containing (12). Rearrange rows as specified there, and we are done.

It remains to show that Steps 3, 6, 9, and 12 above could actually be carried out, by showing that there was sufficient divisibility by x . This will follow from Theorem 4.2.

Definition 4.1. Let $\Delta(0) = 3$ and $\Delta(1) = 2$. For $i \geq 2$, let $b(i)$ denote the largest integer $\leq i$ of the form $2^t - 1$ or $3 \cdot 2^t - 1$, and let $\Delta(i) = i - b(i)$.

For example, the values of $\Delta(i)$ for $2 \leq i \leq 17$ are as in the following table.

i	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
$\Delta(i)$	0	0	1	0	1	0	1	2	3	0	1	2	3	0	1	2

Theorem 4.2. Let $\nu(i, j)$ denote the exponent of x in $P_{i,j}$ at any stage of the reduction from the end of Step 1 to the end of Step 14. Then

- If $19 \leq j \leq 22$ and $0 \leq i \leq j - 8$, then $\nu(i, j) \geq 22 - j + \Delta(i)$.
- If $39 \leq j \leq 46$ and $0 \leq i \leq j - 16$, then $\nu(i, j) \geq 46 - j + \Delta(i)$.
- If $79 \leq j \leq 94$ and $0 \leq i \leq j - 32$, then $\nu(i, j) \geq 94 - j + \Delta(i)$.

Since this applies to any stage of the reduction, it says that all x -exponents in these columns are nonnegative at the end of Steps 3, 6, 9, and 12, which means that there was enough x -divisibility to perform the step. The divisibility of other columns in rows 2, 5, 11, and 23 at Steps 3, 6, 9, and 12 is easily checked, mostly following from proportionality.

Proof. We give the proof for $79 \leq j \leq 94$. The proof for the smaller ranges is basically the same. The proof is by induction on j . By Theorem 2.1(e) shifted, at the outset $\nu(32, 79) = 16$, while $\nu(i, 79) = \infty$ for $i \neq 32$ and $i \leq 47$. If $j \geq 80$, we assume the result is known for $j - 1$. With the rearranging and shifting, we start with, for $i \geq 2$,

$$\nu(i, j) = \begin{cases} \nu_E(i - 1, j - 1) & i \notin \{2, 3, 6, 12, 24, 48\} \\ \nu_E(\frac{1}{2}i - 1, j - 1) & i \in \{6, 12, 24, 48\} \\ \nu_E(i - 2, j - 1) & i \in \{2, 3\}, \end{cases}$$

where $\nu_E(-, -)$ refers to the value of ν at the end of Step 14. By the induction hypothesis, this is

$$\geq \begin{cases} 94 - j + 1 + \Delta(i - 1) & i \notin \{2, 3, 6, 12, 24, 48\} \\ 94 - j + 1 + \Delta(\frac{1}{2}i - 1) = 94 - j + 1 + \Delta(i - 1) & i \in \{6, 12, 24, 48\} \\ 94 - j + 1 + 5 - i & i \in \{2, 3\}. \end{cases}$$

Let $\mu(i, j)$ denote a lower bound for $\nu(i, j) - (94 - j)$. At the outset, we have, for all $i \geq 4$ and $j \geq 79$,

$$\mu(i, j) \geq \Delta(i - 1) + 1,$$

while $\mu(2, j) \geq 4$ and $\mu(3, j) \geq 3$.

We will go through the steps of the reduction and see how μ changes. We can dispense with j as part of the notation. We will now call it $\mu(i)$. To emphasize that μ is changing, we will let μ_k denote the value of μ after Step k . We have $\mu_0(i) \geq \Delta(i - 1) + 1$ for $i \geq 4$, $\mu_0(2) \geq 4$ and $\mu_0(3) \geq 3$. Although it is possible that actual divisibility could increase after a step (by having terms of smallest exponent cancel), our lower bounds, being just bounds, cannot see this. Thus we always have $\mu_{k+1}(i) \leq \mu_k(i)$, so we wish to prove that $\mu_{14}(i) \geq \Delta(i)$.

Step 1 sets

$$(17) \quad \mu_1(1) \geq \min(\mu_0(5), \mu_0(9), \mu_0(17), \mu_0(33), \mu_0(10), \\ \mu_0(18), \mu_0(34), \mu_0(20), \mu_0(36), \mu_0(40)) = 2$$

and $\mu_1(0) = \mu_1(1) + 1 \geq 3$. Of course, $\mu_1(i) = \mu_0(i)$ for $i > 1$, since Step 1 is only changing R_0 and R_1 . In asserting (17), it is relevant that the various \tilde{R}_i which affect R_1 do not include $i = 2^t$ or $3 \cdot 2^t$, since those are the only i for which $\mu_0(i) = 1$.

Step 2 sets

$$\mu_2(2) \geq \min(\mu_1(2), \mu_1(4), \mu_1(8), \mu_1(16), \mu_1(32)) \geq 1.$$

Other rows that affect R_2 would contribute exponents at least this large. Step 3 subtracts 1 from $\mu_2(2)$, so now $\mu_3(2) \geq 0$. Step 4 sets

$$\mu_4(3) \geq \min(\mu_3(3), \mu_3(2)) \geq 0.$$

We have $\mu_4(5) = \mu_0(5) \geq 2$. Step 5 does not change this estimate, i.e., $\mu_5(5) \geq 2$, because at Step 5, R_5 is not affected by any of the rows, $i = 2^t$ with $t \geq 1$ or $i = 3 \cdot 2^t$ with $t \geq 0$, for which $\mu_4(i) < 2$. This is due to the fact that, for these values of i , \tilde{R}_k is 0 in C_{i-1} throughout the reduction for all $k \geq 2$. Step 6 subtracts 2 from $\mu_5(5)$, so now $\mu_6(5) \geq 0$.

For Step 7, we need to know the x -exponents of the entries in C_{10} at this stage of the reduction. These exponents in row i will be 3, 2, 0, 0, 1, 1, 0 for $i = 0, 1, 2, 3, 4, 6$, and 7. These can be seen in the table at

the end of Step 4, or by noting that the entries in rows 4, 6, and 7 will be unchanged from their values in Table 5, while R_1 got x^2 from \tilde{R}_5 at Step 1, R_2 got x from \tilde{R}_4 at Step 2, then changed to x^0 at Step 3, while R_3 then got x^0 at Step 4. For these values of i , we obtain that $\mu_7(i)$ is \geq the minimum of $\mu_6(i)$ and the exponent listed above. It turns out that the only change is $\mu_7(7) \geq 0$. Our bounds now for i from 0 to 7 are 3, 2, 0, 0, 1, 0, 1, 0.

Since $\mu_7(11) \geq 4$ and $\mu_7(i) \geq 4$ for $19 \leq i \leq 23$ and $35 \leq i \leq 47$, we obtain $\mu_8(11) \geq 4$, and then $\mu_9(11) \geq 0$. For Step 10, we need to know exponent bounds in C_{22} at this stage of the reduction, because it is these multiples of R_{11} that are being subtracted from the row in question. For $i < 15$, they will be the same as the μ -values that we are computing here, because the same steps apply. However, we have $\mu_{10}(15) = 0$ due to the $4p_3(x^4)$ -entry in $P_{15,22}$. Our exponent bounds $\mu_{10}(i)$ now for i from 0 to 15 are 3, 2, 0, 0, 1, 0, 1, 0, 1, 2, 3, 0, 1, 2, 3, 0.

Since $\mu_{10}(23) \geq 8$ and $\mu_{10}(i) \geq 8$ for $39 \leq i \leq 47$, we obtain $\mu_{11}(23) \geq 8$, and then $\mu_{12}(23) \geq 0$. For Step 13, we need to know exponent bounds in C_{46} at this stage of the reduction, because it is these multiples of R_{23} that are being subtracted from the row in question. For $i < 31$, they will be the same as the μ -values that we are computing here, because the same steps apply. However, we have $\mu_{13}(31) = 0$ due to the $2p_3(x^8)$ -entry in $P_{31,46}$.

In Step 14, we obtain $\mu_{14}(47) = 0$, with no other changes to μ . Our final values for $\mu_{14}(i)$ are 0 for $i = 2, 3, 5, 7, 11, 15, 23, 31$, and 47, and increasing in increments of 1 from one of these to the next. This equals $\Delta(i)$, as claimed. \square

5 Proof of Theorem 2.2

In this section, we prove Theorem 2.2 by defining a sequence of matrices N_0, \dots, N_{e-1} at various stages of the reduction, and then show that N_s reduces to N_{s+1} . After its rows are rearranged, N_{e-1} will become the matrix described in Theorem 2.2. We explain in Theorem 5.2 how N_0 is obtained from N_{e-1} . Comparing with the case $e = 6$, N_0 through N_4 are the matrix after Steps 1, 4, 7, 10, and 13, respectively, while N_5 is the matrix at the end of Step 14 except that $P_{95,94}$ has not yet been cleared out.

In the following, $\lg(-)$ denotes $\lceil \log_2(-) \rceil$, and $\delta_{i,j}$ is the usual Kronecker symbol. We continue to suppress e from the notation.

Definition 5.1. For $0 \leq s \leq e-1$, N_s is a matrix with rows numbered from 0 to $3 \cdot 2^{e-1} - 1$, and columns from 0 to $3 \cdot 2^{e-1} - 2$ satisfying

a. Its leading entries are

- 2^e in $(0,0)$ and 2^{e-1} in $(1,1)$;
- for $0 \leq k \leq e-1$, 2^{e-k} in $(i, i-1)$ for

$$3 \cdot 2^{k-1} \leq i \leq 3 \cdot 2^k - \begin{cases} 2 & k \leq s-1 \\ 1 & k \geq s; \end{cases}$$

- for $1 \leq \ell \leq s$, $2^{e-\ell-1}$ in $(3 \cdot 2^{\ell-1} - 1, 3 \cdot 2^\ell - 2)$.

b. For $2 \leq \ell + 2 \leq t \leq e$, it has

- $2^{e-t} x^{2^\ell} p_{2^t-\ell-2}(x^{2^\ell})$ in $(2^{\ell+1} + m - 1 - \delta_{\ell+m,0}, 2^t + 2^\ell - 2 + m)$ for

$$\begin{cases} 0 \leq m \leq 2^\ell - 1 & \ell < s \\ 0 \leq m \leq 2^\ell + 0 & \ell = s \\ 1 \leq m \leq 2^\ell + 0 & \ell > s; \end{cases}$$

- $2^{e-t} p_{2^t-\ell-1}(x^{2^\ell})$ in $(3 \cdot 2^\ell + m - 1, 2^t + 2^\ell - 2 + m)$ for

$$1 \leq m \leq 2^\ell + \begin{cases} -1 & \ell < s \\ 0 & \ell \geq s, \end{cases}$$

and in $(\lceil 3 \cdot 2^{\ell-1} \rceil - 1, 2^t + 2^\ell - 2)$ if $\ell \leq s$.

c. Except for the leading entries described in (a),

- all entries in C_j are 0 for $3 \cdot 2^k - 1 \leq j \leq 4 \cdot 2^k - 2$, $k \geq 0$, as are those in $C_{3 \cdot 2^k - 2}$ if $k < s$, while the only additional nonzero entry in $C_{3 \cdot 2^s - 2}$ is 2^{e-s-1} in row $3 \cdot 2^{s-1} - 1$;
- if $t \geq 2$ and $j = 2^t + d$ with $-1 \leq d \leq 2^{t-1} - 2$, then $P_{i,j} = 0$ if $i \geq d + 2^{\lg(d+1.5)+1} + 2$;
- if $k \geq 0$ and $i = 3 \cdot 2^k - 1$, then $P_{i,j} = 0$ for $i \leq j < 2i$.

d. For $3 \leq t < u \leq e$, $2^{t-2} - 1 \leq d \leq 2^{t-1} - 2$, and $i \leq d + 2^{t-1}$,

$$P_{i,2^u+d} = \frac{1}{2^{u-t}} p_{2^{u-t+1}-1}(x^{2^{t-1}}) P_{i,2^t+d}.$$

This is also true for $d = 2^{t-2} - 2$ if $s \geq t - 2$, except in row $3 \cdot 2^{t-3} - 1$.

e. If $2 \leq t \leq e - 1$, $3 \cdot 2^t - 2^{t-1} - 1 \leq j \leq 3 \cdot 2^t - 2$, and $i \leq j - 2^t$, then $P_{i,j}$ is divisible by x^ν with $\nu = 3 \cdot 2^t - 2 - j + \eta_s(i)$, where

$$\eta_0(i) = \begin{cases} 3 - i & 0 \leq i \leq 1 \\ 6 - i & 2 \leq i \leq 3 \\ i - c(i) + 1 & i \geq 4, \end{cases}$$

with $c(i)$ the largest integer $\leq i$ of the form 2^v or $3 \cdot 2^v$, and

$$\eta_s(i) = \begin{cases} 0 & \text{if } i + 1 = 3 \cdot 2^v \text{ or } 4 \cdot 2^v \text{ for } 0 \leq v < s \\ \eta_0(i) & \text{otherwise.} \end{cases}$$

Theorem 2.2 is an immediate consequence of the following result, together with the discussion preceding Step 0 of Section 4.

Theorem 5.2. *Let N_s denote the matrices of Definition 5.1.*

1. After subtracting $2R_{3 \cdot 2^{e-2}-1}$ from $R_{3 \cdot 2^{e-1}-1}$ and then rearranging rows, N_{e-1} satisfies the properties of Theorem 2.2. Call this rearranged matrix Q . The rearranging is that for $i = 3 \cdot 2^t - 1$ with $0 \leq t \leq e - 2$, R_i moves to position $2i$, while for other values of $i > 2$, R_i moves to position $i - 1$.
2. Delete the last column of Q , precede this by a column of 0's, and precede this by the following two rows.

$$\begin{array}{c|cccccccc} & 0 & 1 & 2 & 3 & & 2^e - 1 & 2^e & & 3 \cdot 2^{e-1} - 2 \\ \hline 0 & 2^e & \binom{2^e}{2}x & \binom{2^e}{3}x^2 & \binom{2^e}{4}x^3 & \cdots & x^{2^e-1} & 0 & \cdots & 0 \\ 1 & 2^e & \binom{2^e}{2} & \binom{2^e}{3} & \binom{2^e}{4} & \cdots & 1 & 0 & \cdots & 0 \end{array}$$

Then perform the 2^e -analogues of Steps 0 and 1 of Section 4. The result is the matrix N_0 .

3. For $0 \leq s \leq e - 2$, the matrix N_s reduces to N_{s+1} .

Proof. Part 1 is straightforward but tedious and mostly omitted. As an example of the comparison, the final case of the second \bullet of Definition 5.1(b), after rearranging and changing t to T , says

$$P_{3 \cdot 2^\ell - 2, 2^T + 2^\ell - 2} = 2^{e-T} p_{2^T - \ell - 1}(x^{2^\ell}).$$

With $\ell = s$ and $T = s + t + 1$, this becomes the case $i = 3 \cdot 2^s - 2$ of Theorem 2.2(i).

Next we address Part 2. After shifting and performing Step 0 of Section 4, we will have the 2^e analogue of Table 5, in which we recall that odd factors were not written. It is easy but tedious to verify that everything except rows 0 and 1 will be as stated for N_0 . For example, the first \bullet of Definition 5.1(b) with its $s = 0$, and t replaced by T becomes

$$P_{2^{\ell+1} + m - 1, 2^T + 2^\ell - 2 + m} = 2^{e-T} x^{2^\ell} p_{2^T - \ell - 2}(x^{2^\ell}) \text{ for } 1 \leq m \leq 2^\ell$$

for $\ell > 0$. With $\ell = s$ and $t = T - s$, this matches with part ii of Theorem 2.2 shifted 2 down and 1 to the right.

Part e of Definition 5.1 for Part 2 is somewhat delicate. We had $\eta_{e-1}(i) = 0$ for $i = 2, 3, 5, 7, 11, 15, \dots$, i.e. $i = 2^t - 1$ or $3 \cdot 2^t - 1$, with η_{e-1} increasing by 1's between these values of i . The rearranging done in Part 1 puts these 0's in $i = 4, 2, 10, 6, 22, 14, \dots$, i.e. $i = 2^t - 2$ or $3 \cdot 2^t - 2$, with η again increasing by 1's between these values of i . Shifting these down by 2, as is done in Part 2, puts the 0's in 2^t and $3 \cdot 2^t$, starting with $i = 4$, but we add 1 to the η values because of the shift of columns. For example, column 21 had $\nu \geq 1 + \eta$, but this now applies to column 22, where it is interpreted as $0 + (\eta + 1)$. The values of $\eta_0(2)$ and $\eta_0(3)$ are 1 greater than $\eta_{e-1}(0)$ and $\eta_{e-1}(1)$, respectively. These values are all as claimed of $\eta_0(i)$ for $i \geq 2$.

We kill the terms in R_0 and R_1 except for those in columns of the form $2^t - 1$ by the method of Step 1 of Section 4. For example, if j is of the form $3 \cdot 2^t - 1$ or $5 \cdot 2^t - 1$, $t \geq 0$, then the 2-exponent in R_0 and R_1 is 1 greater than that in R_{j+1} , which is a leading entry. We subtract multiples of $2R_{j+1}$ to kill the terms. This brings up multiples of $2\tilde{R}_{j+1}$. If this is nonzero in C_k , the term brought up can be killed by subtracting a multiple of R_{k+1} . This brings up multiples of \tilde{R}_{k+1} . Because columns 11–15, 23–31, etc., i.e. those j satisfying $3 \cdot 2^t - 1 \leq j \leq 4 \cdot 2^t - 1$, are 0, we will not bring up \tilde{R}_i for i from 12–16, 24–32, etc., and these are the

only rows which contain entries which do not satisfy the proportionality and x -divisibility conditions stated in d and e of Definition 5.1, and the only rows that will have $\eta(i) < 2$. Thus we will obtain $\eta_0(1) \geq 2$, and $\eta_0(0) \geq 3$ since R_0 has an extra factor of x as compared to R_1 .

Similar reasoning applies to columns j not of the form $3 \cdot 2^t - 1$ or $5 \cdot 2^t - 1$. If also $j \neq 2^t - 1$, then the 2-exponent in R_0 and R_1 will exceed that in R_{j+1} by more than 1. We can use an even multiple at one of the two steps of the previous paragraph, or can break it up into more steps, which will make the rows eventually brought up have larger values of i , but, either way, we will not be bringing up the bad rows such as 12–16, etc., and so all the properties will be transferred to R_0 and R_1 . Changing the terms $-\binom{2^e}{2^t}$ in C_{2^t-1} to 2^{e-t} is accomplished similarly, using that these differ by a multiple of 2^{e-t+2} , while the entry in $(2^t, 2^t - 1)$ has 2-exponent $e - t + 1$.

There are three steps to the reduction in Part 3, analogous to Steps 5, 6, and 7 in Section 4. Note that the only nonzero entries of N_s in $C_{3 \cdot 2^s - 2}$ are 2^{e-s-1} in $R_{3 \cdot 2^{s-1} - 1}$, and 2^{e-s} in $R_{3 \cdot 2^s - 1}$, and the second nonzero entry in $R_{3 \cdot 2^s - 1}$ is $2^{e-s-2} x^{2^s} p_2(x^{2^s})$ in $C_{3 \cdot 2^{s+1} - 2}$. The first step is to subtract $2R_{3 \cdot 2^{s-1} - 1}$ from $R_{3 \cdot 2^s - 1}$. If $\tilde{R}_{3 \cdot 2^{s-1} - 1}$ has $q \neq 0$ in C_j , then the $-2q$ brought into $R_{3 \cdot 2^s - 1}$ can be killed by adding qR_{j+1} . The net effect is to remove the leading entry of $R_{3 \cdot 2^s - 1}$, making the $2^{e-s-2} x^{2^s} p_2(x^{2^s})$ in $C_{3 \cdot 2^{s+1} - 2}$ its new leading entry, and to bring into this row various $q\tilde{R}_{j+1}$ for which $P_{3 \cdot 2^{s-1} - 1, j} \neq 0$. By (c), such j must satisfy $j > 2^{s+2} + 1$, and then nonzero entries in \tilde{R}_j only occur in columns $> 2^{s+3} + 1$. This extends the first \bullet in (c) to include also $k = s$, which is needed for N_{s+1} .

We must also consider the effect of these changes on $\eta(3 \cdot 2^s - 1)$. We had $\eta_s(3 \cdot 2^s - 1) = \eta_0(3 \cdot 2^s - 1) = 2^s$. It follows from (c) that none of the j 's appearing above can satisfy $3 \cdot 2^t - 1 \leq j \leq 3 \cdot 2^t + 2^s - 3$ or $4 \cdot 2^t - 1 \leq j \leq 4 \cdot 2^t + 2^s - 3$, $t \geq s$, which are the only values having $\eta_s(j+1) < 2^s$. Thus $\eta(3 \cdot 2^s - 1)$ does not change at this step.

The second step divides $R_{3 \cdot 2^s - 1}$ by $x^{2^s} p_2(x^{2^s})$. This can be done because $\eta_s(3 \cdot 2^s - 1) \geq 2^s$. The dividing changes $\eta(3 \cdot 2^s - 1)$ to 0, which is consistent with the claim for $\eta_{s+1}(3 \cdot 2^s - 1)$. This step changes $P_{3 \cdot 2^s - 1, 2^u + 2^{s+1} - 2}$ from $2^{e-u} x^{2^s} p_{2^u - s - 2}(x^{2^s})$ to $2^{e-u} p_{2^u - s - 1 - 1}(x^{2^{s+1}})$ for $u \geq s + 2$. It removes the entry in the first \bullet of (b) with $\ell = s$, $m = 2^s$, $t = u$ and adds the final entry in the second \bullet of (b) with $\ell = s + 1$ and $t = u$. Now $C_{3 \cdot 2^{s+1} - 2}$ has

- 2^{e-s-2} in row $3 \cdot 2^s - 1$;
- $2^{e-s-2}p_3(x^{2^s})$ in row $2^{s+2} - 1$;
- multiples of 2^{e-s-2} in rows 0 through $2^{s+2} - 1$;
- a leading 2^{e-s-1} in row $3 \cdot 2^{s+1} - 1$;
- other entries 0.

Now we subtract multiples of row $3 \cdot 2^s - 1$ from all other rows except row $3 \cdot 2^{s+1} - 1$ to make them 0 in column $3 \cdot 2^{s+1} - 2$. By property (d), this will zero all entries in column $2^u + 2^{s+1} - 2$, $u > s + 2$, except in rows $3 \cdot 2^s - 1$, $2^{s+2} - 1$, and $2^u + 2^{s+1} - 1$. For $u > s + 2$, the entry in $(2^{s+2} - 1, 2^u + 2^{s+1} - 2)$ is changed from $2^{e-u}p_{2^u-s-1}(x^{2^s})$ to

$$\begin{aligned} & 2^{e-u}(p_{2^u-s-1}(x^{2^s}) - p_3(x^{2^s})p_{2^u-s-1-1}(x^{2^{s+1}})) \\ = & -2^{e-u}x^{2^{s+1}}p_{2^u-s-1-2}(x^{2^{s+1}}). \end{aligned}$$

The minus here can be changed to plus by modifying by a multiple of row $2^u + 2^{s+1} - 1$, which will not affect the properties such as (d) and (e). Property (d) will now hold in N_{s+1} for proportionality out of $C_{2^{s+3}+2^{s+1}-2}$, to the extent claimed there. This change removes the entry of the second • of (b) with $\ell = s$, $m = 2^s$, and $t = u$ and replaces it by the entry of the first • with $\ell = s + 1$, $m = 0$, and $t = u$.

Finally we consider the effect of this step on x -divisibility. If j is as in (e) with $t > s + 1$, and $i \leq 2^{s+2} - 1$ and $i \neq 3 \cdot 2^s - 1$, then the new value of $P_{i,j}$ will equal

$$P_{i,j}^{\text{old}} - \frac{P_{i,3 \cdot 2^{s+1}-2}}{2^{e-s-2}} \cdot P_{3 \cdot 2^s-1,j}.$$

The old $P_{i,j}$ is divisible by $x^{3 \cdot 2^t - 2 - j + \eta_s(i)}$. Also, $P_{i,3 \cdot 2^{s+1}-2}$ is divisible by $x^{\eta_s(i)}$ if $i \leq 2^{s+2} - 2$, and by x^0 if $i = 2^{s+2} - 1$. (Note that (e) did not apply in this latter case due to the condition there which here would say $i \leq j - 2^{s+1}$.) We now have $P_{3 \cdot 2^s-1,j}$ divisible by $x^{3 \cdot 2^t - 2 - j}$ since $\eta(3 \cdot 2^s - 1)$ became 0 at the previous substep. Thus the x -divisibility of $P_{i,j}$ does not decrease except when $i = 2^{s+2} - 1$, where it changes to 0, consistent with $\eta_{s+1}(2^{s+2} - 1) = 0$. \square

6 An easily-checked proof for $e \leq 5$

In this section, we give an easily-checked proof of Theorem 1.3 for $e \leq 5$. Its discovery used the reduced form for M_4 described in Section 2, and a **Mathematica** calculation by González for the M_5 analogue. However, checking its validity only requires elementary verifications.

It is proved in [5, Proposition 4.1] that Theorem 1.3 would follow from showing that

$$(18) \quad 2^{e-k} u^{3 \cdot 2^{k-1} - 3} [0, 0] \neq 0 \text{ in } M_e \text{ for } 1 \leq k \leq e.$$

For $e \leq 5$, (18) is an immediate consequence of the following, which is the main result of this section.

Theorem 6.1. *For $e \geq 1$ and $1 \leq k \leq \min(e, 5)$, there is a homomorphism $\phi_{k,e} : M_e \rightarrow \mathbb{Z}/2^{k+e-1}$ sending $2^{e-k} u^{3 \cdot 2^{k-1} - 3} [0, 0]$ nontrivially.*

The homomorphism $\phi_{k,e}$ is nonzero only on the component of M_e in grading $2(3 \cdot 2^{k-1} - 3)$. The component of M_e in grading $2d$ is generated by the same monomials $u^{d-i-j} [i, j]$ for any e , but the relations depend on e . We will give an explicit formula for $\phi_{k,e}(u^{3 \cdot 2^{k-1} - 3 - i - j} [i, j]) \in \mathbb{Z}$ for $i, j \geq 0$, which is independent of e . Thus we usually call it just ϕ_k . We will prove that ϕ_k applied to a relation (3) in M_e is divisible by 2^{k+e-1} . Since part of our formula is $\phi_k(u^{3 \cdot 2^{k-1} - 3} [0, 0]) = 2^{2k-2}$ and hence

$$\phi_k(2^{e-k} u^{3 \cdot 2^{k-1} - 3} [0, 0]) = 2^{k+e-2} \neq 0 \in \mathbb{Z}/2^{k+e-1},$$

Theorem 6.1 will follow. The hope was to see a pattern in the formulas for ϕ_k that might extend to all k , but they seem a bit too delicate for that.

Since the exponent of u in $u^{3 \cdot 2^{k-1} - 3 - i - j} [i, j]$ is determined by k , i , and j , we do not list it. We write $\phi_k(i, j)$ for $\phi_k(u^{3 \cdot 2^{k-1} - 3 - i - j} [i, j])$, and will sometimes omit the subscript k . We have $\phi_1(0, 0) = 1$, and the only relation in grading 0 in M_e is $2^e [0, 0]$, which handles the case $k = 1$.

Here are the lists of values of $\phi_k(i, j)$ when $k = 2$ and $k = 3$.

$$[4 \mid 0, 0 \mid 2, 2, 2 \mid 0, 1, 1, 0],$$

$$[16 \mid 0, 0 \mid 0, 0, 0 \mid 0, 0, 0, 0 \mid 8, 0, 8, 0, 8 \mid 0, 8, 0, 0, 8, 0 \mid 0, 0, 4, 0, 4, 0, 0 \mid 0, 4, 4, 4, 4, 4, 4, 0 \mid 0, 0, 6, 6, 4, 6, 6, 0, 0 \mid 0, 0, 0, -1, -1, -1, -1, 0, 0, 0]$$

Our functions always satisfy $\phi(i, j) = \phi(j, i)$. The first line says that the nonzero values of ϕ_2 are $\phi_2(0, 0) = 4$, $\phi_2(2, 0) = \phi_2(1, 1) = 2$, and $\phi_2(2, 1) = 1$, and their flips. The next pair of lines says, for example, that $\phi_3(0, 0) = 16$ and

$$\phi_3(i, 6 - i) = \begin{cases} 4 & i = 2, 4 \\ 0 & i = 0, 1, 3, 5, 6. \end{cases}$$

Before we list the formulas for ϕ_4 and ϕ_5 , we discuss the verification that $\phi_{3,e} : M_e \rightarrow \mathbb{Z}/2^{e+2}$ is well-defined for all $e \geq 3$. This one is simple enough that it can be (and was) done by hand. We first consider the case $e = 3$. The coefficients $\binom{8}{1}, \dots, \binom{8}{8}$ in (3) are of the form $8, 4\alpha, 8\alpha', 2\alpha'', 8\beta, 4\beta', 8, 1$, where the α 's are $3 \pmod{4}$, and the β 's odd. There are 55 relations after symmetry is taken into account, but only 13 of them contain any term for which $\nu(\binom{8}{\ell+1}\phi(i-\ell, j)) < 5$. The most delicate is the case $i = 5, j = 4$, in which we have

$$\begin{aligned} & 8\phi(5, 4) + 4\alpha\phi(4, 4) + 8\alpha'\phi(3, 4) + 2\alpha''\phi(2, 4) + 8\beta\phi(1, 4) + 4\beta'\phi(0, 4) \\ &= 8 \cdot 1 - 4\alpha \cdot 4 + 8\alpha' \cdot 4 - 2\alpha'' \cdot 4 + 8\beta \cdot 8 + 4\beta' \cdot 8 \\ &\equiv 8 + 16 + 0 - 24 + 0 + 0 \equiv 0 \pmod{32}. \end{aligned}$$

If $e > 3$, then it is as if the binomial coefficients are multiplied by 2^{e-3} . Their odd factors change, but where it matters, the odd factors are still $3 \pmod{4}$. So ϕ applied to each relation is divisible by $2^{e-3} \cdot 32$. Terms with $\binom{2^e}{9}$ and $\binom{2^e}{10}$ also appear, but they are multiplied by $\phi(0, 0)$ or $\phi(0, 1)$, and so yield multiples of 2^{e+2} . This establishes the well-definedness of $\phi_{3,e}$, and that of $\phi_{2,e}$ is much easier.

Next we list values of $\phi_4(i, j)$ in rows of fixed $i + j$ for which there are some nonzero values. We precede the row by the value of $i + j$. For example, the third listed row says that

$$\phi_4(i, 10 - i) = \begin{cases} 32 & i = 2, 8 \\ 0 & i = 0, 1, 3, 4, 5, 6, 7, 9, 10. \end{cases}$$

0 : 64

8 : 32, 0, 0, 0, 32, 0, 0, 0, 32

10 : 0, 0, 32, 0, 0, 0, 0, 32, 0, 0

12 : 0, 0, 0, 0, 16, 0, 0, 0, 16, 0, 0, 0, 0

14 : 0, 0, 16, 0, 16, 0, 16, 0, 16, 0, 16, 0, 0

15 : 0, 0, 0, 0, 16, 16, 0, 0, 16, 16, 0, 0, 0, 0

$$\begin{aligned}
16 &: 0, 0, 0, 0, 8, 0, 8, 0, 16, 0, 8, 0, 8, 0, 0, 0, 0 \\
17 &: 0, 16, 0, 16, 16, 8, 0, 16, -8, -8, 16, 0, 8, 16, 16, 0, 16, 0 \\
18 &: 0, 0, 0, 0, 8, 8, 4, 8, -4, 0, -4, 8, 4, 8, 8, 0, 0, 0, 0 \\
19 &: 0, 8, 8, 0, 0, -4, -4, -4, 4, 8, 8, 4, -4, -4, -4, 0, 0, 8, 8, 0 \\
20 &: 0, 0, -4, -4, -4, 0, -6, -6, -4, 2, 4, 2, -4, -6, -6, 0, -4, -4, -4, 0, 0 \\
21 &: 0, 0, 0, 6, 6, 6, 0, 3, 3, 1, -1, -1, 1, 3, 3, 0, 6, 6, 6, 0, 0, 0.
\end{aligned}$$

These numbers were discovered using Table 1. Because of the way that they were obtained, it better be the case that they send all relations to 0, at least when $e = 4$. The beauty is that despite the hard work that went into obtaining them, once we have them, it is a simple computer check to verify that they work. It is just a matter of reading these numbers $\phi_4(i, j)$ into the computer and then having the computer check that

$$\sum_{\ell=0}^i \binom{16}{\ell+1} \phi_4(i-\ell, j) \equiv 0 \pmod{128} \text{ for } 0 \leq i \leq 21, 0 \leq j \leq 21-i.$$

Now we can prove by induction on e that if $e > 4$, then

$$\sum_{\ell=0}^i \binom{2^e}{\ell+1} \phi_4(i-\ell, j) \equiv 0 \pmod{2^{e+3}} \text{ for } 0 \leq i \leq 21, 0 \leq j \leq 21-i.$$

It is easy to prove that, for $1 < \ell < 2^{e+1}$,

$$(19) \quad \nu\left(\binom{2^{e+1}}{\ell} - 2\binom{2^e}{\ell}\right) = 2e + 1 - \lceil \log_2(\ell - 1) \rceil - \nu(\ell).$$

The induction argument follows from this and the values of $\phi_4(-)$ listed above. Indeed, the induction step requires

$$(20) \quad \nu(\phi_4(i-\ell, j)) \geq \lceil \log_2(\ell) \rceil + \nu(\ell + 1) - 1,$$

and since $i + j \leq 21$, we have $\nu(\phi_4(i-\ell, j)) \geq 1, 2, 3, 4, 5, 6$ if $\ell \geq 1, 2, 4, 6, 10, 14$, respectively, from which (20) follows.

Our treatment for ϕ_5 is similar. Because of the longer lists, we take advantage of symmetry, and only list $\phi_5(i, j)$ for $i \leq j$. As before, we list values of $\phi_5(i, j)$ in rows of fixed $i + j$ for which there are some nonzero

values. We precede the row by the value of $i + j$. If $i + j = 2t + 1$ (resp. $2t$), the last entry listed is $\phi_5(t, t + 1)$ (resp. $\phi_5(t, t)$).

0 : 256
 16 : 128, 0, 0, 0, 0, 0, 0, 0, 128
 20 : 0, 0, 0, 0, 128, 0, 0, 0, 0, 0, 0
 24 : 0, 0, 0, 0, 0, 0, 0, 0, 64, 0, 0, 0, 0
 28 : 0, 0, 0, 0, 64, 0, 0, 0, 64, 0, 0, 0, 64, 0, 0
 30 : 0, 0, 0, 0, 0, 0, 0, 0, 0, 64, 0, 64, 0, 0, 0
 32 : 0, 0, 0, 0, 0, 0, 0, 0, 32, 0, 0, 0, 32, 0, 0, 0, 64
 33 : 0, 0, 0, 0, 0, 0, 0, 0, 64, 0, 0, 0, 0, 0, 64
 34 : 0, 0, 64, 0, 0, 0, 64, 0, 64, 0, -32, 0, 0, 0, 64, 0, 32, 0
 35 : 0, 0, 0, 0, 0, 0, 0, 0, 64, 64, 64, 0, 64, 64, 0, 64, 0
 36 : 0, 0, 0, 0, 0, 0, 0, 0, 32, 0, 32, 0, 16, 0, 32, 0, -16, 0, 0
 37 : 0, 0, 0, 0, 0, 0, 0, 0, 32, 0, 0, 32, 32, 0, 0, 32, 0, 0
 38 : 0, 0, 32, 0, 32, 0, 0, 0, 0, 16, 0, -16, 32, -16, 0, 16, 32, 0, 0
 39 : 0, 0, 0, 0, 0, 32, 32, 0, 0, 0, 0, 32, 16, 16, 0, 0, 16, 16, 32
 40 : 0, 0, 0, 0, 16, 0, 16, 0, 16, 0, 32, 32, 24, 0, 56, 32, 16, 32, 56, 32, 48
 41 : 0, 32, 0, 32, 32, 48, 0, 0, 16, 48, 0, 48, 48, 56, 0, 16, 8, 24, 16, 16, 8
 42 : 0, 0, 0, 0, 16, 16, 8, 16, 8, 16, 40, 0, 40, 8, 36, 8, 28, 16, 52, 8, 28, 48
 43 : 0, 16, 16, 0, 0, 8, 8, 24, 8, 8, 8, 24, 0, 28, 28, 12, 4, 24, 8, 12, 12, 8
 44 : 0, 0, 24, 24, 24, 0, 28, 28, 16, 4, 28, 24, 0, 0, 18, 2, 4, 26, 28, 2, 20, 2, 20
 45 : 0, 0, 0, 12, 12, 12, 0, 10, 10, 14, 14, 4, 8, 10, 0, 15, 15, 5, 3, 15, 9, 1, 15

The computer checks that

$$\sum_{\ell=0}^i \binom{2^e}{\ell+1} \phi_5(i - \ell, j) \equiv 0 \pmod{2^{e+4}} \text{ for } 0 \leq i \leq 45, 0 \leq j \leq 45 - i$$

is true for $e = 5$. It is then proved for all $e \geq 5$ by induction, using (19) as in the previous case.

Donald M. Davis
Department of Mathematics,
 Bethlehem,
 PA 18015, USA,
 dmd1@lehigh.edu

References

- [1] Astey L.; *Geometric dimension of bundles over real projective spaces*, Quart.J. Math. Oxford **31** (1980), 139–155.
- [2] Farber M., *Topological complexity of motion planning*, Discrete Comput. Geom. **29** (2003), 211–221.
- [3] González J., *Connective K-theoretic Euler classes and nonimmersions of 2^k -lens spaces*, J. London Math. Soc. **63** (2001), 247–256.
- [4] González J., *Topological robotics in lens spaces*, Math. Proc. Cambridge Philos. Soc. **139** (2005), 469–485.
- [5] González J.; Velasco M., Wilson W., *Biequivariant maps of spheres and topological complexity of lens spaces*, Comm. Contemp. Math **15** (2013), 33.
- [6] González J.; Zárate L., *BP-theoretic instabilities to the motion-planning problem in 4-torsion lens spaces*, Osaka J. Math **43** (2006), 581–596.